

CROSSOVER

WG FUNGICIDE

ACTIVE CONSTITUENTS: 375 g/kg CYPRODINIL
250 g/kg FLUDIOXONIL

multi-mode fungicidal action



SUPERIOR BOTRYTIS CONTROL

- Unique chemistry & featuring 2 discrete modes of action.
- Rainfast 2 hours after application.
- Systemic & Protectant action.

SureFire

CROSSOVER

WG FUNGICIDE

ACTIVE CONSTITUENTS: 375 g/kg CYPRODINIL
250 g/kg FLUDIOXONIL

SureFire CROSSOVER WG is a selective fungicide for the control of various fungal diseases in Cucumber, Grapes, Green Beans, Green Peas, Leafy Vegetables, Lettuce, Garlic, Strawberries, Alliums, Capsicum, Cut flowers, Nursery stock and Ornamentals, Onions and Pyrethrum.

SureFire CROSSOVER is a water - dispersible granule pre-pack mixture of cyprodinil and fludioxonil. Cyprodinil is a systemic compound that is taken up into the cuticle and waxy layers of leaves and fruit. From there it is distributed and penetrates into other parts of the plant. Fludioxonil is the residual component of the Crossover mixture and stays mainly on the leaf and fruit surfaces.

Key benefits of CROSSOVER use are:

- 2 discrete modes of action from 2 active ingredients
- Systemic & protectant action. Cyprodinil is systemic & Fludioxonil is residual.
- Extensive range of crops on the label.
- Suitable for either dilute or concentrate spray applications.
- Rainfast 2 hours after application when used with a non-ionic surfactant.
- Especially effective for controlling Botrytis & Sclerotinia rot in leafy vegetables.

In order to minimise the likelihood of the development of fungicide resistance, it is recommended that SUREFIRE CROSSOVER be used in a program incorporating all available forms of crop protection. Alternating or mixing fungicides with different modes of actions is a recognised disease resistance management strategy.

Refer to the table to the right for a full description of the label crop claims for SureFire Crossover.



CROP/SITUATION	DISEASE	RATE			
Capsicum (open field only)	Botrytis rot (<i>Botrytis cinerea</i>)	100 g/100 L OR 1 kg/ha	Lettuce	Sclerotinia Rot (Lettuce Drop) <i>Sclerotinia minor</i> and <i>Sclerotinia sclerotiorum</i>	800 g to 1 kg/ha
	Sclerotinia rot (<i>Sclerotinia minor</i> or <i>S. sclerotiorum</i>)				
Cucumber	Grey Mould (<i>Botrytis cinerea</i>)	800 g to 1 kg/ha		Grey Mould (<i>Botrytis cinerea</i>)	
Cut Flowers	Botrytis rot	80 g/100 L or 800 g/ha plus 0.02% non- ionic surfactant		Anthraco- se (<i>Microdochium panattonianum</i>)	1 kg/ha in a spray volume of 250 – 500 L water/ha
Grapes (except grapes grown for dried fruit production)	Grey Mould (Bunch Rot) (<i>Botrytis cinerea</i>)	Dilute Spraying 80 g/100 L Concentrate Spraying 800 g/ha	Nursery stock (non- food) and ornamentals (non- food) including seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses and fruit trees (non- bearing)	Anthraco- se (<i>Colletotrichum spp.</i>) Grey mould, Rhizoctonia rot (<i>Rhizoctonia spp.</i>), Sclerotinia rot (<i>Sclerotinia spp.</i>)	80 g/100 L or 800 g/ha
Green beans	Grey Mould (<i>Botrytis cinerea</i>) Sclerotinia (<i>Sclerotinia minor</i> and <i>Sclerotinia sclerotiorum</i>)	800 g to 1 kg/ha	Onions	Neck rot (<i>Botrytis alli</i>) Suppression: Black mould (<i>Aspergillus niger</i>)	800 g/ha to 1 kg/ha plus a registered rate of non- ionic surfactant
Green peas including Garden peas, Snow peas and Sugar Snap peas	Grey Mould (<i>Botrytis cinerea</i>) Sclerotinia (<i>Sclerotinia minor</i> and <i>Sclerotinia sclerotiorum</i>)	800 g to 1 kg/ha	Alliums including bulb onions, spring onions, shallots and garlic	Black Mould (<i>Aspergillus niger</i>), suppression only Grey Mould (<i>Botrytis cinerea</i>) suppression only	
Leafy Vegetables (including spinach, silver beet, kale, endive, mustard, cress, chard, rocket, Asian leafy greens and Chinese cabbage)	Sclerotinia (<i>Sclerotinia minor</i> and <i>Sclerotinia sclerotiorum</i>)	800 g to 1 kg/ha	Pyrethrum (<i>Tanacetum cinerariifolium</i>) crops	Winter dieback complex (Ray blight; <i>Phoma ligulicola</i> and Tan spot (<i>Microsphaeropsis tanacetii</i>)	1 kg/ha
			Strawberries	Crown and Petiole Rots (<i>Colletotrichum gloeosporioides</i>) Grey Mould (<i>Botrytis cinerea</i>)	80 g/100 L water

Container sizes: Carton quantity:

1 Kg 10

5 Kg 2

Note: A maximum of 2 applications of SureFire Crossover should be applied per season to ensure effective fungicide resistance management. WHP vary between 3 days to 4 weeks for harvest. Refer to the full label for specific information.

GROUP 9 12 FUNGICIDE

APVMA Approval No: 91395 / 132095

© PCT International
Pty Ltd

www.pctrural.com.au