

INVICTUS 500 WG

FUNGICIDE

500 g/Kg TRIFLOXYSTROBIN

**powdery mildew
apple, pear, grape
downy mildew**



SureFire

INVICTUS 500 WG

FUNGICIDE

500 g/Kg TRIFLOXYSTROBIN

SureFire INVICTUS 500WG fungicide is our new strobilurin based fungicide which offers a robust preventative action against black spot disease in apples and pears (see label segment below).

SureFire INVICTUS exhibits mesostemic properties, i.e. having a high affinity for the surface of the plant, allowing it to be readily absorbed into the waxy layer on the leaf and stems, then it penetrates plant tissue by translaminar activity.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Apples Pears	Powdery mildew of apples Protective control of: Apple scab, Pear scab	Dilute spraying 10 g/100 L water Concentrate Spraying Refer to the Mixing/ Application section	This use is subject to a CropLife Australia resistance management strategy. Follow a complete disease management program for scab by rotating with fungicides from unrelated chemical groups according to the CropLife Australia fungicide resistance management guidelines. Surefire Invictus 500 WG should be applied as a block of three treatments at 7 to 10 day intervals over flowering and into early fruit development. Use short spray intervals when disease pressure is high. DO NOT apply more than three applications of Surefire Invictus 500 WG per season (part of the CropLife Australia resistance management strategy). When used over flowering, apply Surefire Invictus 500 WG alone. Apply with a DMI (Group 3) scab fungicide if intervals are longer than 10 days. Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Do not use in equipment that requires rates greater than 50 g/100 L of water (5X).
Grapevines	Powdery mildew, Suppression of downy mildew	Dilute spraying 15 g/100 L water Concentrate Spraying Refer to the Mixing/ Application section.	Apply thoroughly as part of a powdery mildew spray program. The critical crop timing for powdery mildew control is from early flowering to pre bunch closure. DO NOT allow spray intervals in the above program to exceed 14 days. DO NOT apply more than two applications of Surefire Invictus 500 WG per season. Apply by dilute or concentrate spraying equipment. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Do not use in equipment that requires rates greater than 75 g/100 L of water (5X). Do not apply in water volumes less than 250 L/ha. This use is subject to a CropLife Australia resistance management strategy. For further information refer to the CropLife Australia website: www.croplifeaustralia.org.au

Container sizes:

Carton quantity:

1 Kg
5 Kg

5
4

WITHHOLDING PERIODS:

Apples, pears, grapes: **DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION**

APVMA Approval No: 85778/114048

GROUP **11** FUNGICIDE

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SureFire

INVICTUS 500 WG*

FUNGICIDE

ACTIVE CONSTITUENT: 500 g/kg TRIFLOXYSTROBIN

GROUP **11** FUNGICIDE

For the control of certain diseases of apples, grapevines and pears as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Contents: 5 Kg

PCT

HOLDINGS PTY LTD

APVMA Approval No: 85778/114048

*SUREFIRE Invictus 500 WG is a registered trademark of PCT INTERNATIONAL PTY LTD
(PCT Holdings Pty Ltd ABN 11 099 023 962)

5/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pctrural.com.au>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

© **PCT International Pty Ltd**

www.pctrural.com.au