

STELLAR TOP*

FUNGICIDE

200 g/L AZOXYSTROBIN
125g/L DIFENOCONAZOLE

for top control of
powdery mildew
& blight



CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SureFire®

STELLAR TOP

FUNGICIDE

ACTIVE CONSTITUENTS: 200 g/L AZOXYSTROBIN
125 g/L DIFENOCONAZOLE

GROUP **3 11** FUNGICIDE

For the control of various diseases of Carrots,
Potatoes and Tomatoes as per the Directions for Use.

IMPORTANT:
READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

Contents 5 Litres

PCT
HOLDINGS PTY LTD

APVMA Approval No: 86917/117048

*SureFire Stellar Top is a registered trademark of PCT INTERNATIONAL PTY LTD

(PCT Holdings Pty Ltd ABN 11 009 023 962)

5/74 Murdoch Circuit, Acacia Ridge QLD 4110 - <http://pctrural.com.au>

CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

SureFire

STELLAR TOP®

FUNGICIDE

200 g/L AZOXYSTROBIN
125g/L DIFENOCONAZOLE

Surefire STELLAR TOP is a fungicide registered for control of leaf blight & powdery mildew in carrots, and for early & late blight in potatoes and in tomatoes.

Suitable for both ground & aerial application, tho refer to the label for complete information.

Generally rainfast after 2 hours.

Being a dual mode Group 3 –11 fungicide, it offers an excellent method of fungal resistance control.

Surefire STELLAR TOP is a systemic fungicide, ensures rapid uptake and increased crop yields.

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Carrots	Leaf Blight (<i>Alternaria dauci</i> and <i>Cercospora carotae</i>)	500 or 1000 mL/ha	7 days	Use in a protectant program commencing before disease occurs. Application Apply in a sufficient volume of water to achieve thorough coverage of all foliage. Apply the 500 mL rate at 7 day intervals and the 1000 mL rate at 10 to 14 day intervals. Use the higher rates when climatic conditions are humid and mild, which are favourable for disease infection.
	Powdery Mildew (<i>Erysiphe hercleyi</i>)	1000 mL/ha		Use in a protectant program commencing before disease occurs, particularly during weather conditions that favour disease development. Apply at 10 to 14 day intervals in a sufficient volume of water to achieve thorough coverage of all foliage.
CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Potatoes	Early blight (Target spot) (<i>Alternaria solani</i>)	375 - 500 mL/ha	7 days	Apply the higher rates when climatic conditions favour Early blight or Late blight infection and in crops with large canopies. Application SUREFIRE STELLAR TOP may be applied by ground or aerial application equipment in potatoes. Aerial application may be used only for Early blight (Target spot) control. Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the potatoes.
	Late blight (<i>Phytophthora infestans</i>)	625 - 750 mL/ha		
Tomatoes (except greenhouse)	Early blight (Target spot) (<i>Alternaria solani</i>)	500 mL/ha or 50 mL/100 L	3 days	Application Apply in a sufficient volume of water to achieve thorough coverage of all foliage. The volume of water required to achieve this will depend on the stage of growth of the tomatoes and the method of trellising which influences canopy volume. (refer to sections on Dilute and Concentrate Spraying). In the case of dilute spraying (mL/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy. In the case of fully trellised tomatoes at full canopy, application volumes should be increased to 1500 L/ha to achieve these results with high volume spraying. For Late blight and Sclerotinia control use the higher rates when climatic conditions are humid and mild, which are favourable for disease infection.
	Late blight (<i>Phytophthora infestans</i>), Sclerotinia (<i>Sclerotinia minor</i>)	625 - 750 mL/ha or 63 - 75 mL/100 L		
	Powdery Mildew (<i>Leveillula taurica</i> and <i>Oidium lycopersici</i>)	500 or 750 mL/ha or 50 or 75 mL/100 L		

Container sizes: Carton quantity:

1 litre 5

5 litre 4

Note: No more than 6 applications of SureFire Stellar Top should be applied per season on potatoes or 3 for carrots, for effective fungal resistance management. The Product is phytotoxic to certain apple varieties. Refer to the label for specific information.

GROUP **3|11** FUNGICIDE

APVMA Approval No: 86917 / 117048



© PCT International
Pty Ltd

www.pctrural.com.au