

TRIADIMENOL 250 EC

FUNGICIDE

250 g/L TRIADIMENOL



**Get
control
of
powdery
mildew
in
grapes**

SureFire

TRIADIMENOL 250 EC

FUNGICIDE

250 g/L TRIADIMENOL

Surefire Triadimenol is a broad spectrum, systemic, foliar fungicide.

As it is quickly taken up by plants and rapidly translocated with systemic activity, Surefire Triadimenol is less affected by rainfall or irrigation than contact type fungicides.

It is registered for control of powdery mildew & ring spot on many crops (see table below).

Being a Group 3 fungicide, it fits in well with an insect resistance management strategy.

Suitable for dilute or concentrate spraying.

Withholding periods vary between nil for sugar cane to 7 days for grapes.

Crop	Disease	State	Rate	Critical Comments
Brassica vegetables (eg Cabbage Cauliflower Broccoli, Brussels sprouts)	Ring Spot	All States	200 OR 400 mL/ha plus Agridex 200 mL/ha	Apply every 10 days. Start the program immediately following the first two consecutive days after transplanting. Use the higher rate under heavy disease conditions favouring development of the disease.
Cucurbits	Powdery mildew		40 mL/100 L or 400 mL/ha	Apply a non-DMI fungicide such as oxythioquinox on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Surefire Triadimenol 250 EC Fungicide with a non-DMI fungicide such as oxythioquinox on a 5-10 day schedule.
Tomatoes Peppers (Capsicums and Chillies) Eggplants				Apply Surefire Triadimenol 250 EC Fungicide prior to, or at first sign of disease. Best control is achieved when applied as part of a well planned preventative spray program, with early applications before heavy disease pressure develops.
Grapevines			Dilute Spraying 10 mL/100 L or 100 mL/ha Concentrate Spraying Refer to the Application section	This use pattern is subject to a CropLife Resistance Management strategy (www.croplife.org.au) Apply a maximum of 3 sprays per season, thoroughly and only as part of a spray program. Typically: 1. When shoots are 10-20 cm long. 2. Pre flowering. 3. At flowering. 4. After fruit set. 5. Before bunches close.
Pawpaws	Powdery mildew	QLD, NSW, WA only	Dilute Spraying 20 mL/100 L or 200 mL/ha Concentrate Spraying Refer to the Application section	Spray when powdery mildew infection begins or when conditions favour the disease. Repeat application every 14 days while conditions favour the disease. Ensure thorough coverage of foliage particularly in the crown region. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Sugar Cane	Pineapple disease	QLD, NSW only	20 mL/100 L	Apply to setts by dipping or spraying. Ensure thorough wetting of cut ends.

Container sizes: Carton quantity:

1 litre 5
5 litre 4

APVMA Approval No: 84304/116695

GROUP 3 FUNGICIDE



WITHHOLDING PERIODS

HARVEST:

CUCURBITS, EGGPLANT, PEPPERS (CAPSICUM AND CHILLI), TOMATOES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

BRASSICA VEGETABLES (CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS), GRAPEVINES, PAWPAWS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

SUGAR CANE: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

BRASSICA VEGETABLES: DO NOT GRAZE ANY TREATED AREA, OR CUT FOR STOCK FOOD.

© PCT International Pty Ltd

www.pctrural.com.au