

PCT Holdings Pty Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under off-label permits not endorsed by PCT Holdings Pty Ltd, or under abnormal conditions.

STAROX 400*

ACTIVE CONSTITUENT: 400 g/L FLUOROXYPR as the methyl heptyl ester
SOLVENTS: 300 g/L HYDROCARBON LIQUID
100 g/L N-METHYL PYRROLIDONE

For the control of a wide range of broadleaf weeds in Fallow, Lucerne, Maize, Millet, Pastures, Popples, Sorghum, Sugar Cane, Sweet Corn and Winter Cereals. Also for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, forests, pastures and rights-of-way, as specified in the Directions for Use.



Table 1: Woody weeds in agricultural non-crop areas and rights-of-way, commercial and industrial areas, forests and pastures

HIGH VOLUME APPLICATION: Dilute product with water				3 STATE	water ml/100 L of	CRITICAL COMMENTS
WEEDS CONTROLLED	GROWTH STAGE	Application	GENERAL INSTRUCTIONS			
Barbier burr Negundo burr	Seedlings and young plants up to 40 cm high	NSW, NT, Qld WA only	150			
Black-backed (Common blackberry)	Seedlings and young plants up to 40 cm flowering	NSW, Qld only				
Mimosa paste	Apply from mid to late flowering	NT, WA only				
Common serotine plant Beylania bush	Seedlings and young plants up to flowering	Qld, NSW, WA only	250			
Blackberry Bohara clover	Seedlings and young plants up to 30 cm	Qld, WA only				
Cattail (yellow flower) (Typha angustifolia) (T. monoculosa)	Seedlings and young plants up to 30 cm	NSW, Qld only				
Redberry Cotton weed	Up to 15 cm high flowering					
Cripping banana	Seedlings and young plants up to 30 cm flowering					
Common Docks (Rumex spp.)	Seedlings and young plants up to 30 cm					
Heath scent	Seedlings and young plants up to flowering					
Honey locust	Seedlings and young plants up to 2 m high					
Small flowered mistle (Marrubium) (Marrubium)	Seedlings and young plants up to flowering					
Yellowflower Dwarf's clew	Seedlings and young plants up to flowering					
Canina	High seedlings and young plants up to 1.2 m high		500			

spraying and planting will be determined by the most residual product, i.e. the product with the longest plant-back period.

GROUP	HERBICIDE
-------	-----------

[illegible]

Susceptible crops include but are not limited to clovers, cotton, fruit, trees, lupins, ornamentals, peas, pine trees, potatoes, navy beans, safflower, straw, trunks, soybeans, sunflower, tomatoes, vegetables and vines.

STROX and STROX-400 HERBICIDE can be damaging to susceptible crops during application. Grasses are normally unaffected by STROX and STROX-400 HERBICIDE and are quickly after treatment. Transitory damage can occur on some species, particularly those that spread by stolons such as chard grass (*Cynodon dactylon*), Kikuyu grass and carpet grass (*Axonopus* sp.).

DO NOT allow spray to drift onto susceptible crops, shrubs, trees and *Pinus* spp. DO NOT allow spray to drift onto susceptible crops, shrubs, trees and *Pinus* spp. expected to cause spray to drift onto nearby susceptible plants.

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

Poisonous plants may become more palatable after spraying.

DO NOT allow stock to re-enter paddocks containing treated poisonous plants until the plants have died down.

DO NOT contaminate streams, rivers or waterways with the chemical or used container. Alongside waterways, treat only noxious weeds and poisonous plants.

RESTRAINTS: DO NOT apply to plants which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.

Thorough coverage of both foliage and stems, to the point of runoff, is essential for high volume applications (see GENERAL INSTRUCTIONS; application methods WOODY WEED SITUATIONS section).
DO NOT spray if rain is likely to occur within one hour.

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers and/or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If approved waste management facility is not available bury the empty container in a secure location. DO NOT use for any other purpose. Use up for this purpose clear of walkways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Do not tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter

After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of PCT Holdings Pty Ltd.

Wear protective equipment (see **SAFETY DIRECTIONS**). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see **STORAGE AND DISPOSAL** section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

Additional information is listed in the Safety Data Sheet which can be obtained from your supplier.

Old rough bark will require more spray than smooth or young thin bark.

Apply the recommended mixture liberally to the freshly cut stump immediately after cutting. Apply by spraying or painting the cut surface and sides of the stump. Best results are obtained when stems are cut less than 15 cm above the ground.

Use water should be discharged onto a designated disposal area or, if this is available, onto wasteland away from desirable plants and water courses.

Using: After using SUREFIRE STAROX 400 HERBICIDE, empty the tank completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain, and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the tank as above, quarter fill with clean water and circulate through the pump, lines and nozzles. Drain and repeat the rinsing procedure twice.

PROTECTION OF CROPS): Wash the tank and rinse the system as above. Then refilter fill the tank and add an alkali detergent (e.g. liquid SURF, OMO, DRIVE) at 500 g/100 L of water and circulate throughout the system for at least 15 minutes.

in the whole system. Remove filters and nozzles and clean them separately. Finally, clean the system with clean water and allow to drain.

Remove degreaser. Fill the tank with clean water and add an alkali detergent (e.g. liquid SURF[®], 10, QUART) at 50 ml/10 l of water. Shake the sprayer to circulate the washing liquid throughout the sprayer, then spray the solution through the nozzles. Rinse well with clean water to remove the detergent.

clean brushes and containers, spray liberally with degreaser. Hose off with clean water and repeat using detergents as above.

nt-back periods for crops following the application of SUREFIRE STAROX 400 RBICIDE for rates up to 750 mL/ha

	RATE mL/h	190	375	750
CROP			DAYS	
Barley, Wheat		7	7	7
Chickpea, Sunflower,				
Millet, Sorghum				
Soybean				14
Cotton		14	14	28

Note: Before using SUREFIRE STAROX 400 HERBICIDE in tank mixes with other herbicides, check the plant-back information on all product labels. The time between

