# **POISON**

**KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING** 

# SureFire EXCLAIM 44 SG\*

## **INSECTICIDE**

**ACTIVE CONSTITUENT:** 44 g/kg EMAMECTIN present as EMAMECTIN BENZOATE

GROUP 6 INSECTICIDE

For the control of Diamondback Moth (Cabbage Moth) and Cabbage White Butterfly in Brassica Vegetables and Heliothis in Capsicum, Lettuce and Tomatoes.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

# **Contents 500 grams**



HOLDINGS PTY LTD

APVMA Approval No: 86238/115093

\*SUREFIRE Exclaim 44 SG is a registered trademark of PCT INTERNATIONAL PTY LTD (PCT Holdings Pty Ltd ABN 11 099 023 962)

5/74 Murdoch Circuit, Acacia Ridge QLD 4110 • http://pctrural.com.au CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877

CROP	PEST	RATE	CRITICAL COMMENTS
	Diamondback Moth (Cabbage Moth) (Plutella xylostella) and Cabbage White Butterfly (Pieris rapae)	250 to 300 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. Use the lower rate on low to moderate infestations. Add Citowett or Agral <sup>®</sup> at the appropriate label rate. DO NOT make more than applications to any brassica crop. Where more than one crop is grown DO NOT make more than 4 applications of Surefire Exclaim in any one year. Surefire Exclaim should be used according to CropLife insecticide resistance strategies.
	Heliothis (Helicoverpa armigera and Helicoverpa punctigera)	150 to 250 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <b>Helicoverpe</b> ages have hatched. OB NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year D0 NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 yfa during periods of heavy insect pressure or under very hot and sunny conditions. Surefire Exclaims should be used according to the CropLife resistance strategies.
Tomatoes	Heliothis (Helicoverpa armigera, Helicoverpa punctigera)	150 to 250 g/ha or 15 to 25 g/100 L (dilute spraying) for trellised Tomatoes. Concentrate Spraying Refer to the Mixing/ Application section	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the Helicoverape ageg have hatched. DO NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year D0 NOT apply more than 4 sprays per year. Ensure thorough spray coverage, Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and surmy conditions. Trellised Tomatoes: In the case of dilute spraying (g/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha aft full canopy. Surefire Exclaim should be used according to the Crouli fer resistance strategies.

#### NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Brassica Vegetables: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

DO NOT USE TREATED CROP CROP WASTE OR PRODUCE FOR STOCK FOOD

Capsicum, Lettuce, Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION DO NOT ALLOW LIVESTOCK TO GRAZE TREATED CROPS OR

CROP STURBLE

EXPORT TRADE ADVICE

time being allowed between application and harvest, as some export market have either no Maximum Residue Limits (MRLs) or different MRLs to those of Australia. Details of overseas standards and any export interval can be obtained by contacting PCT Holdings Pty Ltd before using this product.

#### GENERAL INSTRUCTIONS

Add the required amount of Surefire Exclaim to a partly filled INSECTICIDE RESISTANCE WARNING spray tank, and then add the remainder of the water.

### APPLICATION

Surefire Exclaim is not systemic making good coverage essential - ensure thorough coverage of the crop.

#### DO NOT apply by aerial application

Brassica Vegetables: Apply Surefire Exclaim in a minimum of

Capsicums, Lettuce, Tomatoes: Ensure thorough coverage of the crop.

#### Mixing/Application - Trellised Tomatoes

Dilute Spraying: Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

Concentrate Spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute

spray volume (see Dilute Spraying) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated

#### in the following way: Example Only

1. Dilute spray volume as determined above: For example 1000 L/

2. Your chosen concentrate spray volume: For example 500 L/ha 3. The concentration factor in this example is: 2 X (i.e. 1000 L ÷

Treated crop commodities destined for export may require extra 4. If the dilute label rate is 15 g/100 L, then the concentrate rate

becomes 2 x 15, that is 30 g/100 L of concentrate spray The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed

as the crop grows For further information on concentrate spraying, users are advised

to consult relevant industry quidelines, undertake appropriate competency training and follow industry Best Practices.

## GROUP 6 INSECTICIDE

For inwwsecticide resistance management, Surefire Exclaim 44 SAFETY DIRECTIONS SG Insecticide is a Group 6 Insecticide. Some naturally occurring Harmful if swallowed, Will irritate the eves and skin, Avoid contact insect biotypes resistant to Surefire Exclaim and other Group 6 with eyes and skin. When opening the container and preparing insecticides may exist through normal genetic variability in any insect population. Repeated use of one particular insecticide or insecticides from the same chemical group can lead to the early development of insect resistance to that insecticide or chemical group. The effectiveness of Surefire Exclaim on resistant individuals could be significantly reduced. Resistant individuals can eventually dominate the insect population if Surefire Exclaim and other Group 6 insecticides are used repeatedly. Surefire Exclaim should not be applied more than two times per season. and do not apply as consecutive applications. Since occurrence of resistant individuals is difficult to detect prior to use, PCT Holdings Pty Ltd accepts no liability for any losses that may is available from PCT Holdings Ptv Ltd on request. Call Customer. result from the failure of Surefire Exclaim to control resistant Service Toll Free on 1800 630 877 or visit our web site at http:// insects. Surefire Exclaim may be subject to specific resistance management strategies. For further information contact your local supplier, PCT Holdings representative or local agricultural department agronomist.

FNVIRONMENT

Re-entry Period: DO NOT allow entry into treated areas for 8 hours or until the spray has dried. If prior entry is necessary, wear chemical resistant gloves. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND CONDITIONS OF SALE

not burn empty containers or product.

PROTECTION OF LIVESTOCK

STORAGE AND DISPOSAL

ventilated area out of direct sunlight.

bees are foraging.

the product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and goggles or face shield. Wash hands after use. After each day's use, wash gloves, goggles or faceshield and contaminated If poisoning occurs contact a doctor or Poisons Information Centre, Phone Australia 13 11 26. SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet which

Dangerous to fish and other aquatic organisms. DO NOT

contaminate streams, rivers or waterways with the chemical or

used containers, DO NOT apply under weather conditions or from

spraying equipment which could be expected to cause spray to

drift onto adjacent areas, particularly wetlands, water bodies or

Dangerous to bees, DO NOT spray any plants in flower while

Store in the closed, original container in a dry, cool, well

Triple-rinse containers before disposal, Add rinsings to spray

tank. Do not dispose of undiluted chemicals on site. If recycling

replace cap and return clean containers to recycler or designated

collection point. If not recycling, break, crush, or puncture and

deliver empty packaging to an approved waste management

facility. If an approved waste management facility is not available.

bury the empty packaging 500 mm below the surface in a

disposal pit specifically marked and set up for this purpose, clear

of waterways, desirable vegetation and tree roots, in compliance

with relevant local, state or territory government regulations. Do

## pctrural.com.au

PCT Holdings Ptv Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with Directions for Use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the of the product, contrary to label instructions or under off-label permits not endorsed by PCT Holdings Pty Ltd or under abnormal