

PYMENTO

INSECTICIDE

500g/Kg PYMETROZINE

Group 9B
..use for
pesticide
rotation.

aphids?



- systemic action.
- translaminar activity.
- Group 9B mode of action.
- registered for aphids in brassicas, cotton, potatoes, and stone fruit.

SureFire

SureFire PYMENTO

INSECTICIDE

500 g/Kg PYMETROZINE

SureFire Pymento Insecticide contains the active ingredient pymetrozine, which has a specific contact and feed effect against aphids.

The insects stop sucking/eating immediately after intake of Pymento, because the salivary pumps of the insects are being blocked by nerve action of the insecticide. Honey dew production is also quickly stopped.

Although the aphids can stay alive for the first few days after treatment, there is no more reproduction and spreading of the populations.

Furthermore, since Pymento has a unique Group 9B mode of action, it is particularly useful for incorporating in your insect resistance management strategy.



Crop	Pest	Rate	WHP	Critical Comments
Brassica vegetables: Broccoli, Brussels sprouts, Cabbage (except Chinese cabbage), Cauliflowers only	Cabbage aphid, Green peach aphid	200 g/ha	14 days	Add a suitable wetting agent at recommended rates. Minimum retreatment interval 14 days.
Cotton	Cotton Aphid (<i>Aphis gossypii</i>)	400 g/ha Add 0.2%v/v organo-silicone surfactant	4 weeks (H)	Apply to a developing aphid population of no greater than 40% infestation and no more than 5 aphids per leaf. Apply prior to row closure on an actively growing crop before honeydew damage has occurred. Thorough spray coverage is essential. SUREFIRE PYMENTO should be used according to the Cotton Industry's Best Management Practices Manual and its associated Spray and Drift Management Plan. PYMENTO is not a knockdown. DO NOT expect to see a significant reduction in aphid numbers for at least 3 days.
Potatoes	Green peach aphid	200 g/ha	Nil	Minimum retreatment interval 14 days.
Stone fruit	Black peach aphid, Green peach aphid	Dilute spray: 20 g/100 L water Concentrate spray: Refer to the Application – Stone Fruit section	4 weeks	Apply at first sign of aphid infestation. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to General Instructions for application information. Dilute spraying: If water volumes exceed 2000 L per hectare, total rate of SUREFIRE PYMENTO must not exceed 400 g per hectare. Concentrate spraying: DO NOT use at rates greater than 80 g per 100 L of water.

Withholding Periods:
BRASSICAS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION
POTATOES: NOT REQUIRED WHEN USED AS DIRECTED
STONE FRUIT: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION
COTTON: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.
 NOTE: REFER TO LABEL FOR SPECIFICS.

Container sizes: 1 Kg
 Carton quantity: 10

APVMA Approval No: 85119/112318

GROUP 9B INSECTICIDE

© **PCT International Pty Ltd**

www.pctrural.com.au