

COUMA

ALL WEATHER BLOCKS RODENTICIDE

0.37 g/Kg COUMATETRALYL

COUMA RODENT CONTROL

- Registered for rat control in macadamia crops
- Low risk of Secondary Poisoning
- Contains Coumatetralyl as the active Ingredient
- Suitable for indoor & outdoor use



SureFire

COUMA

ALL WEATHER BLOCKS RODENTICIDE

0.37 g/Kg COUMATETRALYL

Surefire COUMA All-Weather Blocks rodenticide is our ready-to-use first-generation multi-feed anticoagulant bait.

- ◆ Highly palatable - no bait shyness
- ◆ Contains COUMATETRALYL as the active ingredient
- ◆ Coumatetralyl is metabolized rapidly in dead rodents, decreasing the risk of secondary poisoning.
- ◆ Registered for use in macadamia, sugar cane & pineapple plantations.
- ◆ Contains Bitrex human-taste deterrent.
- ◆ Suitable for indoor & outdoor use.
- ◆ Approved for use in farm buildings, dairies, meatworks & wharf areas.



Pest	Situation	Rate	Critical Comments
Mice and rats	Domestic and industrial buildings and external areas associated with such establishments, wharf areas, around farm buildings, dairies, meatworks (non-product areas)	30-60 g (1 to 2 blocks) per bait site	Place blocks in areas of known rodent activity. DO NOT exceed 3 m between baits for mice, or 9 m for rats. Follow the Baiting Strategies for non-crop situations detailed below.
Rats	Field crops as below:	60 g (2 blocks per bait station)	Follow the Baiting Strategies for in crop situations detailed below. Bait must be placed in bait stations as described in Preparation of Bait Stations below. Baiting must be used as part of a comprehensive rodent control program. Clear gullies, weeds and scrub. Keep headlands well slashed or bare to discourage rats and to expose them to predators.
	Sugar cane		Practise in-crop grass and weed control. Obtain rat population monitoring data from district organisations and bait according to local recommendations. Strategic baiting is best carried out in December and January.
	Macadamias		If not possible to clear orchard boundaries, rehabilitate to rain forest. Monitor rodent damage levels prior to baiting. Concentrate baiting in the outer 3 rows of crop closest to scrubby habitats and when nuts are available.
	Pineapples		Place bait stations at 9 metre intervals, 1 metre inside the block on all sides where rats are entering the block. Bait should be added to bait stations when fruit begins to form and bait replenished until after harvest.

Do not exceed 3m between bait stations for mice (or 9m for rats).

DO NOT graze or feed livestock on treated areas whilst bait is present. DO NOT place baits in any position accessible to children, livestock or domestic pets.

Do not feed the bait to either wild or domestic birds.

APVMA approval No 82217-105225

Pack sizes:

2.4 Kg & 8Kg pails

© **PCT International Pty Ltd**

www.pctrural.com.au