

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS



**BRODI PELLETS**

ACTIVE CONSTITUENT: 0.05 g/kg BRODIFACOU

**KILLS RATS & MICE**

**FAST ACTING - SINGLE FEED, LETHAL DOSE**

**Controls species resistant to warfarin – CONTAINS BITTERING AGENT**

Ready to use pellets for the control of rats and mice in and around agricultural, industrial, commercial and domestic buildings.

**NET CONTENTS 1 kg**



**APVMA Approval No: 67610/138137**

**\*SUREFIRE® BRODI PELLETS is a registered trademark of PCT INTERNATIONAL PTY LTD**

**(PCT Holdings Pty Ltd ABN 11 099 023 962)**

**5/74 Murdoch Circuit, Acacia Ridge QLD 4110 • <http://pctrural.com.au>**

**CUSTOMER SERVICE FREECALL 1800 630 877 EMERGENCY RESPONSE (ALL HOURS) FREECALL 1800 630 877**

## SUREFIRE BRODI PELLETS

### GENERAL INFORMATION:

Surefire Brodi Pellets is a ready to use, single feed lethal dose product for the control of rats and mice in and around agricultural, industrial, commercial and domestic buildings.

### DIRECTIONS FOR USE

#### Restraints:

DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

DO NOT place bait stations in open.

DO NOT apply bait to crops.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around industrial, commercial, public service, agricultural and domestic buildings.	Rats ( <i>Rattus rattus</i> , <i>Rattus norvegicus</i> )  Mice ( <i>Mus musculus</i> )	50-100 g per bait station depending on level of infestation	Set out bait in places frequented by mice and rats but not more than 2 m from buildings.  <b>For Rats:</b> Place 50-100 g of pellets per station (4-6 g will kill a rat) in 1 or 2 locations at intervals of 5-9 m apart in infested areas. <b>For Mice:</b> Place 50 g of pellets per station (less than 1 g will kill a mouse) in several locations at intervals of 2-3 m apart in infested areas.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities.

### BAITING STRATEGY

Remove, as far as practical, all alternative food sources.

Place pellets in bait stations under cover along walls or rafters and in dark sheltered spots where there are signs of rodent activity.

In drains, suspend bait stations from manholes by wire or place in such a manner so that baits will not be washed into water supplies.

Record the number and location of placements for later replenishment and/or removal.

Inspect daily, replacing eaten baits and increasing the number of placements if they are eaten quickly. Most rodent deaths occur 4 to 7 days after consumption of bait but it is necessary to continue the baiting process for at least 2 weeks or longer; until no more baits are taken. Look for dead rats and mice- burn or bury them, do not place in rubbish bins.

Once effective control has been achieved, remove the bait stations.

### GENERAL INSTRUCTIONS

#### PRECAUTION

Keep away from food, food utensils or places where food is prepared or stored.

### PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, streams, rivers, watercourses or drains with the bait or its used container. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations that are accessible to domestic animals, livestock non-target native animals or birds. The product is restricted to use in and around buildings (within 2 m) or enclosed spaces e.g. drains. Ensure uneaten baits are removed and disposed of with empty containers when rodent activity has ceased.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, puncture and bury empty containers and dispose of unused bait in a local authority landfill. If no landfill is available, bury the containers or unused bait below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

DO NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

### SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT inhale dust. DO NOT touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

### FOR EMERGENCIES ONLY, CALL 1800 033 111 (ALL HOURS)

**Surefire Brodi Pellets contains a bittering agent which helps prevent accidental consumption by children.**

**NOTE TO PHYSICIANS AND VETERINARIANS:** Brodifacoum is a long-lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical or appropriate para-medical personnel or veterinarians. If administered intravenously, the injection must be given slowly.

### SDS

For further information refer to the Safety Data Sheet which is available from the supplier.

**LIABILITY:** This product must be used strictly as directed. PCT Holdings Pty Ltd will not be liable for any loss or damage arising from a failure to strictly follow the directions for use.

**APVMA Approval No.** 67610/138137

**D.O.M.**

**Batch No.:**

